

## JEE (ADVANCED)-2024

## PAPER-1

## CHEMISTRY

## SECTION 1 (Maximum Marks : 12)


- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:


Full Marks : +3 If **ONLY** the correct option is chosen;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);








Negative Marks : 1 In all other cases.

- A closed vessel contains 10 g of an ideal gas X at 300 K, which exerts 2 atm pressure. At the same temperature, 80 g of another ideal gas Y is added to it and the pressure becomes 6 atm. The ratio of root mean square velocities of X and Y at 300 K is  
(A)  $2\sqrt{2} : \sqrt{3}$  (B)  $2\sqrt{2} : 1$  (C) 1 : 2 (D) 2 : 1
- At room temperature, disproportionation of an aqueous solution of in situ generated nitrous acid ( $\text{HNO}_2$ ) gives the species  
(A)  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and NO (B)  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and  $\text{NO}_2$   
(C)  $\text{H}_3\text{O}^+$ ,  $\text{NO}^-$  and  $\text{NO}_2$  (D)  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and  $\text{N}_2\text{O}$
- Aspartame, an artificial sweetener, is a dipeptide aspartyl phenylalanine methyl ester. The structure of aspartame is






4. Among the following options, select the option in which each complex in Set-I shows geometrical isomerism and the two complexes in Set-II are ionization isomers of each other.



### SECTION 2 (Maximum Marks : 12)


- This section contains **THREE (03)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:  
 Full Marks : +4 **ONLY** if (all) the correct option(s) is(are) chosen;  
 Partial Marks : +3 If all the four options are correct but **ONLY** three options are chosen;  
 Partial Marks : + 2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;  
 Partial Marks : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;  
 Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);  
 Negative Marks : -2 In all other cases.

5. Among the following the correct statement(s) for electrons in an atom is(are)
- (A) Uncertainty principle rules out the existence of definite paths for electrons.
- (B) The energy of an electron in 2s orbital of an atom is lower than the energy of an electron that is infinitely far away from the nucleus.
- (C) According to Bohr's model, the most negative energy value for an electron is given by  $n=1$ , which corresponds to the most stable orbit.
- (D) According to Bohr's model, the magnitude of velocity of electrons increases with increase in values of  $n$ .
6. Reaction of iso-propylbenzene with  $O_2$  followed by the treatment with  $H_3O^+$  forms phenol and a by-product **P**. Reaction of **P** with 3 equivalents of  $Cl_2$  gives compound **Q**. Treatment of **Q** with  $Ca(OH)_2$  produces compound **R** and calcium salt **S**.
- The correct statement(s) regarding **P**, **Q**, **R** and **S** is(are)
- (A) Reaction of **P** with **R** in the presence of  $KOH$  followed by acidification gives
- 
- (B) Reaction of **R** with  $O_2$  in the presence of light gives phosgene gas
- (C) **Q** reacts with aqueous  $NaOH$  to produce  $Cl_3CCH_2OH$  and  $Cl_3CCOONa$
- (D) **S** on heating gives **P**
7. The option(s) in which at least three molecules follow Octet Rule is(are)
- (A)  $CO_2$ ,  $C_2H_4$ ,  $NO$  and  $HCl$
- (B)  $NO_2$ ,  $O_3$ ,  $HCl$  and  $H_2SO_4$
- (C)  $BCl_3$ ,  $NO$ ,  $NO_2$  and  $H_2SO_4$
- (D)  $CO_2$ ,  $BCl_3$ ,  $O_3$  and  $C_2H_4$

## SECTION 3 (Maximum Marks : 24)

- This section contains **SIX (06)** questions.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:  
Full Marks : +4 If **ONLY** the correct integer is entered;  
Zero Marks : 0 In all other cases.


8. Consider the following volume-temperature ( $V-T$ ) diagram for the expansion of 5 moles of an ideal monoatomic gas.





Considering only  $P-V$  work is involved, the total change in enthalpy (in Joule) for the transformation of state in the sequence  $X \rightarrow Y \rightarrow Z$  is

[Use the given data: Molar heat capacity of the gas for the given temperature range,  $C_{V,m} = 12 \text{ J K}^{-1} \text{ mol}^{-1}$  and gas constant,  $R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$ ]

9. Consider the following reaction,



which follows the mechanism given below:



The order of the reaction is \_\_\_\_\_?


10. Complete reaction of acetaldehyde with excess formaldehyde, upon heating with conc.  $\text{NaOH}$  solution, gives **P** and **Q**. Compound **P** does not give Tollens' test, whereas **Q** on acidification gives positive Tollens' test. Treatment of **P** with excess cyclohexanone in the presence of catalytic amount of *p*-toluenesulfonic acid (PTSA) gives product **R**.

Sum of the number of methylene groups ( $-\text{CH}_2-$ ) and oxygen atoms in **R** is \_\_\_\_\_.

11. Among  $\text{V}(\text{CO})_6$ ,  $\text{Cr}(\text{CO})_5$ ,  $\text{Cu}(\text{CO})_3$ ,  $\text{Mn}(\text{CO})_5$ ,  $\text{Fe}(\text{CO})_5$ ,  $[\text{Co}(\text{CO})_3]^{3-}$ ,  $[\text{Cr}(\text{CO})_4]^{4-}$ , and  $\text{Ir}(\text{CO})_3$ , the total number of species isoelectronic with  $\text{Ni}(\text{CO})_4$  is

[Given atomic number :  $\text{V} = 23$ ,  $\text{Cr} = 24$ ,  $\text{Mn} = 25$ ,  $\text{Fe} = 26$ ,  $\text{Co} = 27$ ,  $\text{Ni} = 28$ ,  $\text{Cu} = 29$ ,  $\text{Ir} = 77$ ]


12. In the following reaction sequence, the major product **P** is formed.



Glycerol reacts completely with excess **P** in the presence of an acid catalyst to form **Q**. Reaction of **Q** with excess **NaOH** followed by the treatment with **CaCl<sub>2</sub>** yields **Ca**-soap **R**, quantitatively. Starting with one mole of **Q**, the amount of **R** produced in gram is

[Given, atomic weight: H = 1, C = 12, N = 14, O = 16, Na = 23, Cl = 35, Ca = 40]

13. Among the following complexes, the total number of diamagnetic species is \_\_\_\_\_.



[Given, atomic number: Mn = 25, Fe = 26, Co = 27; en = H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>]

#### SECTION 4 (Maximum Marks : 12)

- This section contains **FOUR (04)** Matching List Sets.
- Each set has **ONE** Multiple Choice Question.
- Each set has **TWO** lists: **List-I** and **List-II**.
- List-I** has **Four** entries (P), (Q), (R) and (S) and **List-II** has **Five** entries (1), (2), (3), (4) and (5).
- FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks : +3 **ONLY** if the option corresponding to the correct combination is chosen;

Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);

Negative Marks : 1 In all other cases.






14. In a conductometric titration, small volume of titrant of higher concentration is added stepwise to a larger volume of titrate of much lower concentration, and the conductance is measured after each addition.

The limiting ionic conductivity ( $\Lambda_0$ ) values (in  $\text{mSm}^2 \text{mol}^{-1}$ ) for different ions in aqueous solutions are given below:

Ions	Ag <sup>+</sup>	K <sup>+</sup>	Na <sup>+</sup>	H <sup>+</sup>	NO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	OH <sup>-</sup>	CH <sub>3</sub> COO <sup>-</sup>
$\Lambda_0$	6.2	7.4	5.0	35.0	7.2	7.6	16.0	19.9	4.1

For different combinations of titrates and titrants given in List-I, the graphs of 'conductance' versus 'volume of titrant' are given in List-II.

Match each entry in List-I with the appropriate entry in List-II and choose the correct option.

	List-I		List-II
(P)	Titrate: KCl Titrant: $\text{AgNO}_3$	(1)	
(Q)	Titrate: $\text{AgNO}_3$ Titrant: KCl	(2)	
(R)	Titrate: NaOH Titrant: HCl	(3)	
(S)	Titrate: NaOH Titrant: $\text{CH}_3\text{COOH}$	(4)	
		(5)	

(A) P-4, Q-3, R-2, S-5

(B) P-2, Q-4, R-3, S-1

(C) P-3, Q-4, R-2, S-5

(D) P-4, Q-3, R-2, S-1

15. Based on VSEPR model, match the xenon compounds given in List-I with the corresponding geometries and the number of lone pairs on xenon given in List-II and choose the correct option.

	List-I		List-II
(P)	$\text{XeF}_2$	(1)	Trigonal bipyramidal and two lone pair of electrons
(Q)	$\text{XeF}_4$	(2)	Tetrahedral and one lone pair of electrons
(R)	$\text{XeO}_3$	(3)	Octahedral and two lone pair of electrons
(S)	$\text{XeO}_3\text{F}_2$	(4)	Trigonal bipyramidal and no lone pair of electrons
		(5)	Trigonal bipyramidal and three lone pair of electrons

(A) P-5, Q-2, R-3, S-1










(B) P-5, Q-3, R-2, S-4

(C) P-4, Q-3, R-2, S-1

(D) P-4, Q-2, R-5, S-3












16. List-I contains various reaction sequences and List-II contains the possible products. Match each entry in List-I with the appropriate entry in List-II and choose the correct option.

	List-I		List-II
(P)	 $\xrightarrow{\begin{array}{l} \text{i) O}_3, \text{Zn} \\ \text{ii) aq. NaOH}, \Delta \\ \text{iii) ethylene glycol, PTSA} \\ \text{iv) a) BH}_3, \text{b) H}_2\text{O}_2, \text{NaOH} \\ \text{v) H}_3\text{O}^+ \\ \text{vi) NaBH}_4 \end{array}}$	(1)	
(Q)	 $\xrightarrow{\begin{array}{l} \text{i) O}_3, \text{Zn} \\ \text{ii) aq. NaOH}, \Delta \\ \text{iii) ethylene glycol, PTSA} \\ \text{iv) a) BH}_3, \text{b) H}_2\text{O}_2, \text{NaOH} \\ \text{v) H}_3\text{O}^+ \\ \text{vi) NaBH}_4 \end{array}}$	(2)	
(R)	 $\xrightarrow{\begin{array}{l} \text{i) ethylene glycol, PTSA} \\ \text{ii) a) Hg(OAc)}_2, \text{H}_2\text{O}, \text{b) NaBH}_4 \\ \text{iii) H}_3\text{O}^+ \\ \text{iv) NaBH}_4 \end{array}}$	(3)	
(S)	 $\xrightarrow{\begin{array}{l} \text{i) ethylene glycol, PTSA} \\ \text{ii) a) BH}_3, \text{b) H}_2\text{O}_2, \text{NaOH} \\ \text{iii) H}_3\text{O}^+ \\ \text{iv) NaBH}_4 \end{array}}$	(4)	
		(5)	

- (A) P-3, Q-5, R-4, S-1  
 (B) P-3, Q-2, R-4, S-1  
 (C) P-3, Q-5, R-1, S-4  
 (D) P-5, Q-2, R-4, S-1

17. List-I contains various reaction sequences and List-II contains different phenolic compounds. Match each entry in List-I with the appropriate entry in List-II and choose the correct option.

List-I	List-II
(P)  $\xrightarrow[\text{(ii) Conc. HNO}_3]{\text{(i) molten NaOH, H}_3\text{O}^+}$	(1) 
(Q)  $\xrightarrow[\text{(v) Conc. HNO}_2/\text{Conc. H}_2\text{SO}_4]{\text{(i) Conc. HNO}_3/\text{Conc. H}_2\text{SO}_4, \text{(ii) Sn/HCl, (iii) NaNO}_2/\text{HCl, 0-5}^\circ\text{C, (iv) H}_2\text{O}}$	(2) 
(R)  $\xrightarrow[\text{(iii) H}_3\text{O}^+, \Delta]{\text{(i) Conc. H}_2\text{SO}_4, \text{(ii) Conc. HNO}_3}$	(3) 
(S)  $\xrightarrow[\text{(vi) H}_2\text{O}]{\text{(i) (a) KMnO}_4/\text{KOH, } \Delta; \text{(b) H}_3\text{O}^+, \text{(ii) Conc. HNO}_3/\text{Conc. H}_2\text{SO}_4, \Delta, \text{(iii) (a) SOCl}_2, \text{(b) NH}_3, \text{(iv) Br}_2, \text{NaOH, (v) NaNO}_2/\text{HCl, 0-5}^\circ\text{C}}$	(4) 
	(5) 

(A) P-2, Q-3, R-4, S-5

(B) P-2, Q-3, R-5, S-1

(C) P-3, Q-5, R-4, S-1

(D) P-3, Q-2, R-5, S-4

## ANSWER KEY

- |           |         |         |         |              |              |
|-----------|---------|---------|---------|--------------|--------------|
| 1. (D)    | 2. (A)  | 3. (B)  | 4. (C)  | 5. (A, B, C) | 6. (A, B, D) |
| 7. (A, D) | 8. 8120 | 9. 3    | 10. 18  | 11. 1        | 12. 909      |
| 13. 1     | 14. (C) | 15. (B) | 16. (A) | 17. (C)      |              |